

# Paschen-Back effect in the CrH molecule

New diagnostic tool for measuring magnetic fields in ultra-cool dwarfs

O. Kuzmychov   S.V. Berdyugina

Warsaw  
May 08, 2012



# Today's program

Why we study CrH

How we study CrH


Simulations with STOPRO

Individual lines

0 – 0 band

Outlook

# Why CrH? Fig. from <http://www.stsci.edu/~inr/ldwarf1.html>



# Objectives

- ▶ linelist for CrH,
- ▶ line strengths (in the absence/presence of a mag. field),
- ▶ simulation of a CrH spectra (Stokes I, Q and V),
- ▶ observations.

# Hamiltonian for ${}^6\Sigma$ el. state

$$\hat{H} = \underbrace{\hat{H}_{\text{int}} + \hat{H}_{\text{rot}} + \hat{H}_{\text{so}} + \hat{H}_{\text{sr}} + \dots}_{\text{Ram et al. 1993}} + \underbrace{\hat{H}_H}_{\text{Berdyugina et al. 2005}}$$

- ▶ eigenvalues,
- ▶ eigenvectors.

# Intensity

$$I_{n''}^{n'} \propto A_{n''}^{n'} \propto \sum \left| R_{n''}^{n'} \right|^2,$$

where

$$\sum_{M' M''} \left| R_{n''}^{n'} \right|^2 = f_e \times q_{v' v''} \times \underbrace{\sum_{M' M''} q_{\Omega' \Omega'' J' J'' M' M''}^2}_{\text{line strength}}$$

# Amplitudes

$$q_{\Omega' \Omega'' J' J'' M' M''}^1 = q_{\Omega' \Omega'' J' J''} \times q_{J' J'' M' M''}$$

$$q_{\Omega'_i \Omega''_j J'_k J''_l}^{(b)} = \sum_{\Omega'_i} \sum_{\Omega''_j} C_{\Omega'_i J'_k}^* q_{\Omega'_i \Omega''_j J'_k J''_l}^{(a)} C_{\Omega''_j J''_l}$$

$$q_{\Omega'_k \Omega''_l J'_k J''_l M'_{km} M''_{ln}}^{pb} = \sum_{M'_k} \sum_{M''_l} C_{J'_k M'_{km}}^* q_{\Omega'_k \Omega''_l J'_k J''_l}^{(b)} q_{J'_k J''_l M'_{km} M''_{ln}} C_{J''_l M''_{ln}}$$

Indices:

$$i, j = -5/2, -3/2, \dots, 5/2$$

case (a)

$$k, l = -5/2, -3/2, \dots, 5/2$$

case (b)

$$m, n = -J, -J + 1, \dots, J$$

both cases

---

<sup>1</sup>Schadee 1978

# Recipe for simulations with STOPRO<sup>2</sup>



- ▶ linelist
  - ▶ energies of transitions
  - ▶ strengths of transitions
- ▶ spectroscopic data:  
damping constants, FC factor, ...
- ▶ atmospheric model ( $T = 2500$  K,  $\log g = 4.5$ )
- ▶ strength of the mag. field
- ▶ orientation of the mag. field ( $\chi = 0^\circ$ ,  $\gamma = 45^\circ$ )

---

<sup>2</sup>Solanki 1987 and Solanki et al. 1992





# Zeeman regime





$$\Delta E = g\mu_0 M H$$



# Paschen-Back regime





# Paschen-Back regime





# Paschen-Back regime




# Paschen-Back regime




# Paschen-Back regime



## crh00\_05kG\_chi=0\_gam=45




## crh00\_1kG\_chi=0\_gam=45







## crh00\_3kG\_chi=0\_gam=45



## crh00\_6kG\_chi=0\_gam=45



crh00\_10kG\_chi=0\_gam=45



# Summary & Outlook

- ▶ promising potential of the CrH,
- ▶ observing campaign is pursued,
- ▶ if success  $\Rightarrow$  first mag. field maps for L-dwarfs.